



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Therefore the projectile will fall without the circle, if $\sin\phi$ is less than $\frac{1}{2}(1+\cos\theta)^{\frac{1}{2}} - \frac{1}{2}(1-\cos\theta)^{\frac{1}{2}}$; but will fall within if $\sin\phi$ is greater than $\frac{1}{2}(1+\cos\theta)^{\frac{1}{2}} + \frac{1}{2}(1-\cos\theta)^{\frac{1}{2}}$.

If all possible directions are equally probable, the chance of the projectile falling within the circle is $1 - (1 - \cos\theta)^{\frac{1}{2}} = 1 - \sqrt{2} \sin\frac{1}{2}\theta$.

Hence the required chance is

$$p = \frac{\int_0^{\frac{1}{2}\pi} (1 - \sqrt{2} \sin\frac{1}{2}\theta) d\theta}{\int_0^{\pi} d\theta} = \frac{1}{2} - (2/\pi)(\sqrt{2} - 1).$$

III. Solution by GEORGE LILLEY, Ph. D., Professor of Mathematics, University of Oregon, Eugene, Ore.

The diameter of the field is v^2/g , and the range must not exceed $(v^2/g)\cos\phi$. The elevation may vary from 0° to θ_1 , and from $\frac{1}{2}\pi - \theta_1$ to $\frac{1}{2}\pi$ for each value of ϕ , where θ_1 is determined by $\sin 2\theta_1 = \cos\phi$, ϕ the azimuth of the projectile measured from the diameter, θ the elevation of the gun, v the velocity of projection, and g the intensity of gravity.

The surface-element of the enveloping hemisphere whose radius is R or v^2/g is $R^2 \cos\theta d\theta d\phi$.

Using only one-half the field and one-fourth of the sphere the required chance is

$$\frac{R^2 \int_0^{\frac{1}{2}\pi} \int_0^{\theta_1} \cos\theta d\theta d\phi + R^2 \int_0^{\frac{1}{2}\pi} \int_{\frac{1}{2}\pi - \theta_1}^{\frac{1}{2}\pi} \cos\theta d\theta d\phi}{\pi R^2}$$

$$\text{or } \frac{1}{\pi} \int_0^{\frac{1}{2}\pi} (1 - \sqrt{2} \sin\frac{1}{2}\phi) d\phi, \text{ or } \frac{1}{2} - \frac{2}{\pi}(\sqrt{2} - 1).$$

Solved in a similar manner by L. C. WALKER. Solved with different results by HENRY HEATON, and P. H. PHILBRICK.

75. Proposed by F. P. MATZ, M. Sc., Ph. D., Professor of Mathematics and Astronomy, Irving College, Mechanicsburg, Pa.

Find the mean area of all plane rectilinear right triangles having a constant perimeter p .

I. Solution by G. B. M. ZERR, A. M., Ph. D., Professor of Mathematics and Science, Chester High School, Chester, Pa.

Let x and y be the base and altitude, respectively.

Then $x + y + \sqrt{(x^2 + y^2)} = p$, the perimeter (1).

$$\therefore x = \frac{p^2 - 2py}{2(p - y)}. \quad \therefore \text{Area} = \frac{p(p - 2y)y}{4(p - y)}.$$

The limits of y are $y=0$ to $y=x$. When $y=x$ in (1) we get $y=p/(2+\sqrt{2})=y'$.

$$\begin{aligned}\therefore \Delta &= \frac{p}{4} \frac{\int_0^{y'} \frac{(p-2y)y}{p-y} dy}{\int_0^{y'} dy} = \frac{2+\sqrt{2}}{4} \int_0^y \left(2y+p-\frac{p^2}{p-y}\right) dy \\ &= \frac{2+\sqrt{2}}{4} \left[\frac{1}{(2+\sqrt{2})^2} + \frac{1}{2+\sqrt{2}} + \log \left(\frac{1+\sqrt{2}}{2+\sqrt{2}} \right) \right] p^2 \\ &= \frac{p^2}{4} \left[\frac{1}{2+\sqrt{2}} + 1 + (2+\sqrt{2}) \log \left(\frac{1}{\sqrt{2}} \right) \right] \\ &= \frac{p^2}{8} [4 - \sqrt{2} - (2+\sqrt{2}) \log 2].\end{aligned}$$

II. Solution by J. SCHEFFER, A. M., Hagerstown, Md., and L. C. WALKER, Associate Professor of Mathematics, Leland Stanford University, Palo Alto, Cal.

Denote the base of the right triangle by x , and the perpendicular by y .

Then the hypotenuse is $p-x-y$, and from $(p-x-y)^2=a^2+y^2$, we obtain

$$y = \frac{p^2-2px}{2(p-x)}. \quad \therefore \text{The area} = \frac{p}{4} \frac{(p-2x)x}{p-x}, \text{ and the average area is}$$

$$\begin{aligned}\frac{p}{4} \int_0^{\frac{1}{2}p} \frac{p(p-2x)x}{p-x} dx &\div \int_0^{\frac{1}{2}p} dx = \frac{1}{2} \int_0^{\frac{1}{2}p} \frac{p(p-2x)x}{p-x} dx = \frac{1}{2} \int_0^{\frac{1}{2}p} \left(2x+p-\frac{p^2}{p-x}\right) dx \\ &= \frac{1}{2} p^2 (3-4\log 2).\end{aligned}$$

NOTE.—The difference in the results of these two solutions is due to the different limits assumed for the variable. It seems that the proper limits to assume for the variable x is 0 and $\frac{1}{2}p$, for in this way, and in this way only, do we get the totality of triangles according to our assumed law of distribution of the triangles. The law of distribution tacitly assumed in both solutions is that the number of triangles is proportional to the base of the triangle. Had some other law of distribution been assumed, different results from either of the above would have been obtained. The problem is stated in the indefinite form because the law of distribution is not stated. See Dr. E. H. Moore's Note on Mean Values, Vol. II, page 303, of THE AMERICAN MATHEMATICAL MONTHLY. ED. F.